

Cerebral cortical neuron diversity and development at single-cell resolution

Matthew B Johnson^{1,2,3} and Christopher A Walsh^{1,2,3,4,5,6}



Over a century of efforts to categorize the astonishing diversity of cortical neurons has relied on criteria of morphology, electrophysiology, ontology, and the expression of a few transcripts and proteins. The rapid development of single-cell RNA sequencing (scRNA-seq) adds genome-wide gene expression patterns to this list of criteria, and promises to reveal new insights into the transitions that establish neuronal identity during development, differentiation, activity, and disease. Comparing single neuron data to reference atlases constructed from hundreds of thousands of single-cell transcriptomes will be critical to understanding these transitions and the molecular mechanisms that drive them. We review early efforts, and discuss future challenges and opportunities, in applying scRNA-seq to the elucidation of neuronal subtypes and their development.

Addresses

¹ Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA, USA

² Manton Center for Orphan Disease Research, Boston Children's Hospital, Boston, MA, USA

³ Howard Hughes Medical Institute, Boston Children's Hospital, Boston, MA, USA

⁴ Department of Pediatrics, Harvard Medical School, Boston, MA, USA

⁵ Department of Neurology, Harvard Medical School, Boston, MA, USA

⁶ Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

Corresponding author: Walsh, Christopher A
(christopher.walsh@childrens.harvard.edu)

Current Opinion in Neurobiology 2017, 42:9–16

This review comes from a themed issue on **Developmental neuroscience**

Edited by **Paola Arlotta** and **Pierre Vanderhaegen**

<http://dx.doi.org/10.1016/j.conb.2016.11.001>

0959-4388/© 2016 Elsevier Ltd. All rights reserved.

Introduction

The classification of cell types in the cerebral cortex has challenged the greatest minds in the history of neuroscience, and so perhaps it is no surprise that we do not quite have it figured out yet. Ramon y Cajal and other early histologists described the two major cortical neuronal types — those with long, distantly projecting axons and those with short, locally projecting

axons — and documented their many morphological variations [1]. Brodmann, Campbell, Vogt and others used the distribution of morphological types to subdivide the cortex into cytoarchitectonic areas which we now understand have important functional correlates [2–4]. Yet, whereas classical neuroscientists reached consensus around the neuronal types in brain structures like the cerebellum over a century ago, the effort to develop a comprehensive neuronal ‘parts list’ for the cortex has lagged. Electrophysiological and circuit analyses arrived in the mid-twentieth century with new tools and the idea that morphological and functional classes of neurons might somehow correspond, though the labor-intensive nature of combining electrophysiology and morphology has limited the ability to integrate form and function. The revolution in molecular biology of the late 20th century allowed an integration of developmental lineage, inferred from the expression of a few marker genes [5], yet still it is not clear whether these criteria can define a clean, non-overlapping ‘periodic table’ of cortical neuronal types, or whether instead the classification of cortical neurons is inherently less precise than in other brain areas, with a mix of some sharply defined classes and other, fuzzier categories [6,7]. This review will focus on the relevance of single-cell transcriptomics to the classification of cortical neuron subtypes by genome-wide gene expression, and explore the unique perspective afforded by scRNA-seq on the dynamic processes of cortical neurogenesis and differentiation.

Transcriptomic classification of neuronal cell types

Pioneering single-cell microarray and qRT-PCR studies elucidated progenitor and neuronal subtypes in the mouse brainstem [8], olfactory system [9], retina [10,11], inner ear [12], and embryonic cortex [13,14], as well as developing human and ferret cortex [15]. Now, single-cell RNA-seq has opened the floodgates for deep transcriptomic analysis of CNS cell types [16,17[•],18–26] (see also recent review by [27]). Although some early scRNA-seq studies have tested specific hypotheses — for example, confirming the ‘one neuron-one receptor rule,’ that each individual primary olfactory neuron expresses one and only one olfactory receptor gene [28–31] — most have aimed to elaborate on the molecular identities of classically defined neuronal types, discover new types, and begin to establish definitive brain cell type taxonomies (Table 1). These studies employ a generalizable two-stage approach to scRNA-seq-based cell type classification. In the first stage of analysis, single-cell transcriptomes are grouped through

Table 1

Summary of experimental methods and main results for selected scRNA-seq studies classifying cell types in the mammalian brain. UMI, unique molecular identifiers; ERCC, spike-in synthetic control RNAs.

Reference	Cell selection and isolation method	cDNA type	Number of cells used (total cells sequenced)	Source of cells assayed	Sequencing depth (avg or median reads per cell)	Numbers of genes detected (average per cell and/or cumulative) and used for classification	Cell types identified/ classified	Other notes
Kodama [8]	Manual dissection and cell picking	qRT-PCR (3-Prime end)	167 (208)	Mouse medial vestibular nucleus	N/A	59 hand-picked genes	6 neuronal types: 3 excitatory, 3 inhibitory	Further subdivisions likely, but classification correlates with known morphological and functional subtypes
Saraiva [29]	FACS selection followed by Fluidigm C1	Full-length (SMARTer) with ERCC	21 (58)	Mouse olfactory sensory neurons	4.4 Million	4717 detected per cell; 13 582 total; 509 genes found to be differentially expressed between individual neurons	18 known cell types, one confirmed new cell type	Confirmed the 'one-neuron-one-receptor' hypothesis
Uoskin [17**]	Manual dissection and automated cell picking	5-Prime (STRT) with UMI	622 (799)	Mouse dorsal root ganglion	1.1 Million	3574 ± 2010 detected per cell; 12 750 used to ID initial 4 neuronal subgroups & non-neuronal cells; 11 658 used for final iterative clustering of 11 subtypes	11 neuronal subtypes	68 'outliers or unresolved ID' (8.5% of all cells sequenced)
Darmanis [35]	Fluidigm C1	Full-length (SMARTer)	466 (482)	Human adult temporal lobe and fetal cortex	2.8 Million	~4000 detected per cell	7 neuronal types: 2 excitatory, 5 inhibitory; one fetal progenitor class	16 (3.3%) cells excluded for low reads (<400k)
Tasic [33**]	Manual dissection followed by single-cell FACS into microtiter plates	Full-length (SMARTer) with ERCC	1679 (1739)	Mouse (8 wks) primary visual cortex	8.7 Million	7278 detected per cell; 13 878 used for classification	49 'core' cell types: 19 excitatory, 23 inhibitory, 7 non-neuronal; plus 'intermediate' cells w/mixed identity between two or more 'core' types	255 (15.2%) good cells are of 'intermediate' neuronal subtype
Zeisel [32**]	Manual dissection ± FACS selection followed by Fluidigm C1	5-Prime (STRT) with UMI	3005 (3315)	Mouse (3-5 wks) somatosensory cortex & hippocampus	0.5 Million	~4500 detected per cell; 15k total detected; top 5000 by variance used for classification	47 cell types: 7 excitatory neuron, 16 interneuron, 2 astrocyte, 6 oligodendrocyte, and 2 immune classes	310 outlier/poor quality cells (9.4%)

Table 1 (Continued)

Reference	Cell selection and isolation method	cDNA type	Number of cells used (total cells sequenced)	Source of cells assayed	Sequencing depth (avg or median reads per cell)	Numbers of genes detected (average per cell and/or cumulative) and used for classification	Cell types identified/classified	Other notes
Lake [34**]	FACS selection for NeuN+ nuclei followed by Fluidigm C1	Full-length (SMARTer) with ERCC	3227 (4488)	Human cerebral cortex, six Brodmann areas	8.3 Million	6159 detected per nucleus; 16 242 used for clustering and classification	16 neuronal types: 8 excitatory, 8 inhibitory	28% samples failed initial QC, including ~10% empty samples and ~18% doublets 2% did not map well from the projection set onto the training set; a further 7% did not cluster well
Macosko [45]	Droplets	3-Prime with UMI	13 155 training set; 44 808 total classified (49 300)	Mouse retina	13 000	Training set cells, >900 genes detected; PCA performed on 384 most variable genes; cell type clusters had to have at least 50 cells and 10 differentially expressed genes	39 cell types: 33 neuronal, 6 non-neuronal	

a combination of dimensionality reduction and hierarchical clustering, with varying degrees of iteration and supervision. In the second stage, the resulting cell groups are contrasted against each other to identify differentially expressed marker genes. This approach has so far met with great success in marrying transcriptomic classifications to known cell types. As sample sizes expand and studies proliferate, new methods to standardize these taxonomies and map other data modalities onto transcriptomic cell types will be critical.

The cerebral cortex: the ultimate cell type diversity challenge

Three of the largest scRNA-seq studies of neuronal identity published to date have, fittingly, tackled the most heterogeneous brain region, the cerebral cortex [32**,33**,34**]. In the first such study, unbiased sampling of mouse primary somatosensory and hippocampal cortex identified many non-neuronal cell types in addition to seven excitatory and 16 inhibitory neuronal types, corresponding well to existing layer-defined and marker-defined classes [32**]. Acknowledging that an unbiased sampling captures few cells from rare populations, the authors also oversampled a subtype of interneurons by FACS isolation, enabling them to find a novel subtype of PAX6+ neurogliaform cells in layer I, nicely validated by immunohistochemistry and electrophysiology [32**].

In contrast, selection of known neuronal populations by microdissection and FACS purification from reporter mice yielded a more diverse taxonomy of 42 neuron classes [33**], reassuringly concordant with the selected laminar and marker-based populations, but also further subdividing many of these classes into putative novel subpopulations. In addition, these authors' classification algorithm is relatively lenient with regard to cell type ambiguity, assigning ~15% of cells an 'intermediate' identity between two neuronal classes. In the most extreme case, one third of layer IV neurons were classified as intermediate between two of the three proposed layer IV subtypes. What remains to be determined is the extent to which these intermediate cells reflect algorithmic 'oversplitting,' or subdivision of cell types based on transcriptional variability that is in fact stochastic or state-related rather than subtype identity-dependent. Further studies will be necessary to differentiate state versus trait transcriptional signals in these and any other proposed novel neuronal types.

Adult human brain presents particular challenges for single-cell studies, given the highly myelinated and dense extracellular milieu of the tissue and the typical storage conditions of postmortem samples, which make clean dissociation of intact whole cells difficult [35]. There are, however, reliable and reproducible protocols for isolating single neuronal nuclei from frozen postmortem human brain [36,37]. Fortunately, the nucleus contains

a significant amount of messenger RNA, and several studies have now demonstrated single-nucleus RNA sequencing [38–40]. Lake *et al.* have taken this approach for scRNA-seq-based cell type classification in the human cerebral cortex, identifying 16 neuronal subtypes — 8 excitatory and 8 inhibitory [34**]. Most intriguingly, these authors also detected differences between neocortical areas, for example, between the layer IV transcriptomic subtypes present in frontal versus occipital cortex.

Merging transcriptomic taxonomy with other data types

Given that traditional methods of cell type classification are much lower throughput than scRNA-seq, they are likely to be increasingly guided by single-cell transcriptomic taxonomies, through the ability to collect the transcriptome of a single cell that has already been characterized by another method and map that transcriptome onto large scRNA-seq reference data sets. Several groups have recently demonstrated the collection and sequencing of RNA from cells following electrophysiological recordings [41*,42,43]. In one study, 45 CCK⁺ interneurons in cortical layer I were first characterized by patch-clamp recording and classified into five subtypes based purely on their electrophysiological properties [41*]. Each cell's contents were then aspirated into the patch pipette and the RNA sequenced. The 45 single-cell transcriptomes were then mapped onto 16 interneuron subtypes defined previously [32**]. Interestingly, none of the five electrophysiological subtypes corresponded to a single transcriptomic subtype, with cells from each group mapping to two, three, or even four different scRNA-seq-defined interneuron classes. Similar discord was observed for excitatory neurons.

Discrepancy between electrophysiological and transcriptomic cell type could arise trivially from the technical noise inherent in scRNA-seq, or from differences in experimental conditions. However, it is notable that electrophysiology seems to provide greater unity of CCK⁺ interneuron subtypes compared to scRNA-seq, again suggesting that the algorithm applied to the larger scRNA-seq data set [32**] may have over-split some cell types. Alternatively, the methods used to map one taxonomy onto the other may improve with inclusion of additional *a priori* knowledge; about a third of the

Patch-seq interneurons assayed in somatosensory cortex mapped to subtypes predominantly found in hippocampus in the prior scRNA-seq study. The electrophysiological properties of Patch-seq cells were also highly correlated with their expression of 24 of 167 genes encoding relevant proteins — channels, pumps, receptors, so on. It would be informative to cluster the larger scRNA-seq sample using the same 167-gene panel and map the Patch-seq interneurons onto the resulting taxonomy, which should improve the correspondence between the two data modalities. Indeed, overlapping but distinct sets of genes are likely to determine distinct neuronal properties, including morphology, electrophysiology, and connectivity, as well as developmental processes, like migration, that interact with these features. Although several studies have found a nearly 1:1 correspondence between morphology and electrophysiology for layer I interneurons [42,44], such correspondence is strikingly lacking for interneurons in the rest of the cortical layers [44], supporting the partial independence of these properties. It is likely that classifications based on the summation of all of these transcriptional signals will yield neither unambiguous subtypes nor clear correspondence to individual cellular properties, and the challenge rather will be to deconvolve single-cell transcriptional profiles into the distinct signatures that correspond to each domain of neuronal properties—electrophysiology, location, dendritic arborization, axonal projection, so on.

Droplets and the advantages of 10× higher throughput

Although the methods employed by the vast majority of scRNA-seq studies to date — manual or automated cell picking [17**,28,31]; single-cell flow cytometry [33**]; and/or microfluidics (i.e., the Fluidigm C1 system) [19,29,30,32**,34**,35] — may capture as much as 25% of each cell's mRNA, they are limited in throughput, and by the cost-per-cell for library construction and sequencing (Table 2). An alternative approach involves encapsulating single cells in nanoliter-volume droplets and performing lysis and barcoded reverse transcription within the droplets [26,45,46*]. For comparison, to assay 3300 cortical cells [32**] required weeks of cumulative C1 run time; manual screening of cDNA samples; and dozens of library preparations and sequencing lanes,

Table 2

Pros and cons of current single-cell capture and processing methods.

Cell capture method	Throughput, cells per day	Cost per cell	cDNA output
Manual/automated cell picking	Tens of cells	Tens of dollars	5', 3', or full-length cDNA
Flow cytometry	Hundreds of cells	Dollars	5', 3', or full-length cDNA
Microfluidics	Tens or hundreds	Dollars	full-length cDNA (barcoded)
Droplets	Tens of thousands	Cents	3' tag, already barcoded

* Fluidigm's high-throughput chip is expected to capture up to 800 cells per run, with some level of on-chip barcoding.

whereas a droplet-based study of the mouse retina [46[•]] generated cDNA from 49 300 cells, pooled into seven libraries, over the course of four days — that is, 15 times the number of cells assayed by one fifth the amount of sequencing, and probably less than one tenth the hands-on processing time. The two main drawbacks of current droplet-based methods are (1) only about 10% of a cell's mRNA molecules are captured; and (2) the lack of full-length cDNA generation limits alternative splicing analyses (Table 2), though technological improvements are likely to ameliorate both of these disadvantages in the near future.

Increasing throughput by three orders of magnitude alleviates the pressure to choose between a hypothesis-driven, targeted design that may miss unknown cell types [33^{••}] and an unbiased approach that undersamples minority populations [32^{••}]. In fact, analyzing 49 300 retinal cells first required excluding a large proportion of the rod photoreceptors that, making up two thirds of the retina, masked the transcriptional signatures differentiating other cell types [46[•]]. Following this *in silico* selection step, 39 cell types were defined based on a 'training set' of ~13 000 cells, and the remaining ~36 300 cells were classified by correlation to the training set. Notably, the large sample size enabled identification of known and novel neuronal subtypes as rare as 0.1%. Remarkably however, these subtypes are still far fewer than the 60–100 retinal cell types defined by morphology and physiology [47]. In particular, retinal ganglion cells are known to be highly diverse and yet, because they compose only about 0.5% to 1% of cells in the retina, the current sample of 49 300 cells was insufficient to detect heterogeneity within the ~500 RGCs assayed. Thus, even with the high throughput of droplet-based methods, careful experimental design will be paramount, and a logical expectation is that a complete cell type taxonomy from any tissue that contains rare subpopulations will require a tiered approach of initial unbiased sampling followed by targeted subpopulation studies using known or novel markers to isolate the rarest cells.

Progenitor heterogeneity of the human fetal cortex

Remarkably, while only a single scRNA-seq analysis of mouse embryonic cortex has so far been published [48], a large number of studies have applied single-cell transcriptomics to fetal human cortex [15,35,49[•],50[•],51], motivated by the intriguing morphological heterogeneity of primate cortical neural progenitor cells (NPC) [52–55]. A major contributor to the diversity of primate NPC is the relative abundance of basal or outer radial glia (ORG), which morphologically, functionally, and transcriptionally resemble apical radial glia of the ventricular zone (VZ) but are located in the subventricular zone (SVZ). Two studies employing scRNA-seq to uncover the distinct transcriptional program of ORG employed different selection

methods to enrich for their NPC parent population, and notably arrived at similar results [49[•],50[•]]. By manually microdissecting the VZ/inner SVZ from the outer SVZ before scRNA-seq, Pollen *et al.* were able to subsequently correlate gene expression with germinal zone location and thus identify genes specifically enriched in ORG [49[•]]. At the same time, Thomsen *et al.* developed a protocol for light fixation, permeabilization, and fluorescent immunolabeling compatible with FACS purification of NPC followed by scRNA-seq [50[•]]. Remarkably, both efforts identified several of the same ORG-enriched genes, including *HOPX*, *FAM107A*, and *TNC*. Nevertheless, the sample sizes of these studies are small compared to those on the adult cortex, and much remains to be done. It is important to note that single progenitor transcriptomes are heavily influenced by cell cycle phase, and probably also reflect dramatic changes in fate potential over the course of cortical neurogenesis. Indeed, several ORG-enriched genes were found to be expressed by VZ NPC slightly earlier in development [49[•]]. Altogether, there is a strong case for further developmental studies not only including greater cell numbers but also sampling a wide range of time-points, and applying new analysis methods that can probe the dynamic development and complex lineage relationships of the developing cortex.

Assessing validity and utility of *in vitro* models by scRNA-seq

A key use of scRNA-seq will be to validate *in vitro* models of human brain development and disease by comparing the cell types and developmental dynamics of these models to primary human tissues [51,56,57]. Cerebral organoids are fast becoming a popular model for early human brain development, but neither the variability across iPSC lines or individual organoids nor the correspondence of organoids to *in vivo* brain development is yet fully understood. An scRNA-seq analysis of cerebral organoids at 33–65 days post-differentiation identified both dorsal and ventral telencephalic NPC and neurons, as well as a few cells resembling those of the cortical hem signaling center, an important source of patterning morphogens [51]. Importantly, parallel analyses of organoids and primary human fetal cortex found a remarkable similarity in transcriptional programs of neurogenesis and differentiation, the biggest difference being a paucity of basal NPC in organoids (6% of cells in the organoids vs 34% in 12–13 weeks of gestation human cortex). Future studies will be needed to determine whether this discrepancy reflects the absence or underdevelopment of an important progenitor niche in the organoids, or simply a mismatch in the developmental stages of the organoids and primary tissues compared in this study.

Challenges and opportunities for future studies

Single-cell technologies and data analysis methods continue to improve rapidly, and will be invaluable in creating

a complete census of cell types and lineage relationships in the brain. We foresee future improvements leading to great opportunities in four general areas. Methods to isolate cell populations from non-genetic model species will be of particular importance. The FRISCR method [50*] has great potential, but relies on suitable antibodies and known cell type markers. Promising alternatives include using fluorescent *in situ* hybridization or RT-PCR reactions to sort cells on the abundance of mRNA transcripts [58,59], and merging these methods with droplet-based scRNA-seq will be hugely advantageous. In the model of Patch-seq, protocols are sorely needed for sequencing RNA from single cells previously or concurrently characterized by other methods, for example, physiology, connectivity, developmental lineage, or live imaging. Methods to either maintain [60] or reconstruct [61,62] spatial information in conjunction with scRNA-seq need further development for application to mammalian brain studies. Similarly, emerging methods to sequence DNA and RNA from the same single cell [63,64] will provide critical insights into the lineage relationships between cell types, which are otherwise extremely difficult to assess in human brain. New statistical models have been developed to improve gene expression level estimates and quantify heterogeneity in noisy single-cell data [65,66]. Further development should be aimed at integrating these models with advanced clustering and pseudotime methods. Large-scale developmental studies will require new algorithms for inferring cell type lineages from scRNA-seq data collected at multiple real time points, while normalizing for cell cycle phase. Validation of putative novel subtypes will be paramount, and will require innovative approaches to visualize transcriptional dynamics in single cells over time. The true test of cell type identity is stability, and thus time-lapse live imaging of single-cell transcription will be needed to definitively differentiate transcriptional cell type from cell state.

Conflicts of interest

Nothing declared.

Acknowledgements

M.B.J. is supported by the Paul G. Allen Family Foundation. C.A.W. is an Investigator of the Howard Hughes Medical Institute.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Ramon, Cajal S: *Histologie du système nerveux de l'homme & des vertébrés*. 1909.
 2. Campbell AW: *Histological Studies on the Localisation of Cerebral Function*. 1905.
 3. Brodmann K: *Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf Grund des Zellenbaues*. 1909.
 4. Vogt O: **Die myeloarchitektonische Felderung des menschlichen Stirnhirns**. *J Psychol Neurol* 1910, **15**:221-232.
 5. Lodato S, Arlotta P: **Generating neuronal diversity in the mammalian cerebral cortex**. *Annu Rev Cell Dev Biol* 2015, **31**:699-720.
 6. Masland RH: **Neuronal cell types**. *Curr Biol* 2004, **14**:R497-R500.
 7. Yano K, Subkhankulova T, Livesey FJ, Robinson, Hugh PC: **Electrophysiological and gene expression profiling of neuronal cell types in mammalian neocortex**. *J Physiol (Lond)* 2006, **575**:361-365.
 8. Kodama T, Guerrero S, Shin M, Moghadam S, Faulstich M, Lac du S: **Neuronal classification and marker gene identification via single-cell expression profiling of brainstem vestibular neurons subserving cerebellar learning**. *J Neurosci* 2012, **32**:7819-7831.
 9. Tietjen I, Rihel JM, Cao Y, Koentges G, Zakhary L, Dulac C: **Single-cell transcriptional analysis of neuronal progenitors**. *Neuron* 2003, **38**:161-175.
 10. Trimarchi JM, Stadler MB, Roska B, Billings N, Sun B, Bartch B, Cepko CL: **Molecular heterogeneity of developing retinal ganglion and amacrine cells revealed through single cell gene expression profiling**. *J Comp Neurol* 2007, **502**:1047-1065.
 11. Cherry TJ, Trimarchi JM, Stadler MB, Cepko CL: **Development and diversification of retinal amacrine interneurons at single cell resolution**. *Proc Natl Acad Sci* 2009, **106**:9495-9500.
 12. Durruthy-Durruthy R, Gottlieb A, Hartman BH, Waldhaus J, Laske RD, Altman R, Heller S: **Reconstruction of the mouse otocyst and early neuroblast lineage at single-cell resolution**. *Cell* 2014, **157**:964-978.
 13. Kawaguchi A, Ikawa T, Kasukawa T, Ueda HR, Kurimoto K, Saitou M, Matsuzaki F: **Single-cell gene profiling defines differential progenitor subclasses in mammalian neurogenesis**. *Development* 2008, **135**:3113-3124.
 14. Narayanan G, Poonepalli A, Chen J, Sankaran S, Hariharan S, Yu YH, Robson P, Yang H, Ahmed S: **Single-cell mRNA profiling identifies progenitor subclasses in neurospheres**. *Stem Cells Dev* 2012, **21**:3351-3362.
 15. Johnson MB, Wang PP, Atabay KD, Murphy EA, Doan RN, Hecht JL, Walsh CA: **Single-cell analysis reveals transcriptional heterogeneity of neural progenitors in human cortex**. *Nat Neurosci* 2015, **18**:637-646.
 16. Spaethling JM, Piel D, Dueck H, Buckley PT, Morris JF, Fisher SA, Lee J, Sul J-Y, Kim J, Barfai T et al.: **Serotonergic neuron regulation informed by in vivo single-cell transcriptomics**. *Faseb J* 2014, **28**:771-780.
 17. Usoskin D, Furlan A, Islam S, Abdo H, Lönnerberg P, Lou D, Hjerling-Leffler J, Haeggström J, Kharchenko O, Kharchenko PV et al.: **Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing**. *Nat Neurosci* 2015, **18**:145-153.
- This study exemplifies the analysis approach central to neuron classification by scRNA-seq, with iterative PCA-based partitioning followed by differential expression testing to identify cell types. They recapitulated known developmental, morphological, and physiological sensory neuron classes, and discovered a novel subclassification of C-fiber nociceptors.
18. Li C-L, Li K-C, Wu D, Chen Y, Luo H, Zhao J-R, Wang S-S, Sun M-M, Lu Y-J, Zhong Y-Q et al.: **Somatosensory neuron types identified by high-coverage single-cell RNA-sequencing and functional heterogeneity**. *Cell Res* 2015, **26**:83-102.
 19. Burns JC, Kelly MC, Hoa M, Morell RJ, Kelley MW: **Single-cell RNA-Seq resolves cellular complexity in sensory organs from the neonatal inner ear**. *Nat Commun* 2015, **6**:8557.
 20. Patel AP, Tirosh I, Trombetta JJ, Shalek AK, Gillespie SM, Wakimoto H, Cahill DP, Nahed BV, Curry WT, Martuza RL et al.: **Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma**. *Science* 2014, **344**:1396-1401.
 21. Luo Y, Coskun V, Liang A, Yu J, Cheng L, Ge W, Shi Z, Zhang K, Li C, Cui Y et al.: **Single-cell transcriptome analyses reveal**

- signals to activate dormant neural stem cells. *Cell* 2015, **161**:1175-1186.
22. Shin J, Berg DA, Zhu Y, Shin JY, Song J, Bonaguidi MA, Enikolopov G, Nauen DW, Christian KM, Ming G-L *et al.*: **Single-cell RNA-Seq with waterfall reveals molecular cascades underlying adult neurogenesis.** *Cell Stem Cell* 2015, **17**:360-372.
 23. Gao Y, Wang F, Eisinger BE, Kelnhofer LE, Jobe EM, Zhao X: **Integrative single-cell transcriptomics reveals molecular networks defining neuronal maturation during postnatal neurogenesis.** *Cereb Cortex* 2016 <http://dx.doi.org/10.1093/cercor/bhw040>.
 24. Gökce Ö, Stanley GM, Treutlein B, Neff NF, Camp JG, Malenka RC, Rothwell PE, Fuccillo MV, Südhof TC, Quake SR: **Cellular taxonomy of the mouse striatum as revealed by single-cell RNA-Seq.** *Cell Rep* 2016, **16**:1126-1137.
 25. Marques S, Zeisel A, Codeluppi S, van Bruggen D, Mendanha Falcão A, Xiao L, Li H, Häring M, Hochgerner H, Romanov RA *et al.*: **Oligodendrocyte heterogeneity in the mouse juvenile and adult central nervous system.** *Science* 2016, **352**:1326-1329.
 26. Shekhar K, Lapan SW, Whitney IE, Tran NM, Macosko EZ, Kowalczyk M, Adiconis X, Levin JZ, Nemesh J, Goldman M *et al.*: **Comprehensive classification of retinal bipolar neurons by single-cell transcriptomics.** *Cell* 2016, **166** 1308-1323.e30.
 27. Poulin J-F, Tasic B, Hjerling-Leffler J, Trimarchi JM, Awatramani R: **Disentangling neural cell diversity using single-cell transcriptomics.** *Nat Neurosci* 2016, **19**:1131-1141.
 28. Hanchate NK, Kondoh K, Lu Z, Kuang D, Ye X, Qiu X, Pachter L, Trapnell C, Buck LB: **Single-cell transcriptomics reveals receptor transformations during olfactory neurogenesis.** *Science* 2015, **350**:1251-1255.
 29. Saraiva LR, Ibarra-Soria X, Khan M, Omura M, Scialdone A, Mombaerts P, Marioni JC, Logan DW: **Hierarchical deconstruction of mouse olfactory sensory neurons: from whole mucosa to single-cell RNA-seq.** *Sci Rep* 2015, **5**:18178.
 30. Tan L, Li Q, Xie XS: **Olfactory sensory neurons transiently express multiple olfactory receptors during development.** *Mol Syst Biol* 2015, **11**:844.
 31. Scholz P, Kalbe B, Jansen F, Altmueller J, Becker C, Mohrhardt J, Schreiner B, Gisselmann G, Hatt H, Osterloh S: **Transcriptome analysis of murine olfactory sensory neurons during development using single Cell RNA-Seq.** *Chem Senses* 2016, **41**:313-323.
 32. Zeisel A, Muñoz-Manchado AB, Codeluppi S, Lönnerberg P, La Manno G, Jureus A, Marques S, Munguba H, He L, Betscholtz C *et al.*: **Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq.** *Science* 2015, **347**:1138-1142.
- The first study to apply scRNA-seq to neuron classification in the cerebral cortex highlights the advantages and drawbacks of unbiased sampling of a complex tissue, and uses targeted subsampling to identify a novel interneuron subtype.
33. Tasic B, Menon V, Nguyen TN, Kim T-K, Jarsky T, Yao Z, Levi B, Gray LT, Sorensen SA, Dolbeare T *et al.*: **Adult mouse cortical cell taxonomy revealed by single cell transcriptomics.** *Nat Neurosci* 2016, **19**:335-346.
- This group's unique classification strategy allows for 'intermediate' cell types that exhibit mixed transcriptional signatures between two cortical neuron subtypes. The need for flexible categorization may reflect the complexity of the cerebral cortex and the degree of interplay between 'state' (e.g., highly plastic synaptic connectivity and activity) and 'trait' (cell type).
34. Lake BB, Ai R, Kaeser GE, Salathia NS, Yung YC, Liu R, Wildberg A, Gao D, Fung H-L, Chen S *et al.*: **Neuronal subtypes and diversity revealed by single-nucleus RNA sequencing of the human brain.** *Science* 2016, **352**:1586-1590.
- This study highlights the potential of single-nucleus RNA-sequencing for analyzing difficult-to-dissociate tissues such as postmortem adult human cortex. Analysis of several Brodmann areas also provides an exciting first look at how transcriptomic cell type composition varies across functional domains of the cortex.
35. Darmanis S, Sloan SA, Zhang Y, Enge M, Caneda C, Shuer LM, Hayden Gephart MG, Barres BA, Quake SR: **A survey of human brain transcriptome diversity at the single cell level.** *Proc Natl Acad Sci* 2015, **112**:7285-7290.
 36. Spalding KL, Bhardwaj RD, Buchholz BA, Druid H, Frisén J: **Retrospective birth dating of cells in humans.** *Cell* 2005, **122**:133-143.
 37. Evrony GD, Cai X, Lee E, Hills LB, Elhosary PC, Lehmann HS, Parker JJ, Atabay KD, Gilmore EC, Poduri A *et al.*: **Single-neuron sequencing analysis of L1 retrotransposition and somatic mutation in the human brain.** *Cell* 2012, **151**:483-496.
 38. Krishnaswami SR, Grindberg RV, Novotny M, Venepally P, Lacar B, Bhutani K, Linker SB, Pham S, Erwin JA, Miller JA *et al.*: **Using single nuclei for RNA-seq to capture the transcriptome of postmortem neurons.** *Nat Protoc* 2016, **11**:499-524.
 39. Lacar B, Linker SB, Jaeger BN, Krishnaswami S, Barron J, Kelder M, Parylak S, Paquola A, Venepally P, Novotny M *et al.*: **Nuclear RNA-seq of single neurons reveals molecular signatures of activation.** *Nat Commun* 2016, **7**:11022.
 40. Habib N, Li Y, Heidenreich M, Swiech L, Avraham-Davidi I, Trombetta JJ, Hession C, Zhang F, Regev A: **Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare adult newborn neurons.** *Science* 2016, **353**:925-928.
 41. Fuzik J, Zeisel A, Máté Z, Calvigioni D, Yanagawa Y, Szabó G, Linnarsson S, Harkany T: **Integration of electrophysiological recordings with single-cell RNA-seq data identifies neuronal subtypes.** *Nat Biotechnol* 2016, **34**:175-183.
- This study exemplifies the opportunities and challenges in merging neuronal taxonomies derived from different data modalities, and provides strong evidence that in the cerebral cortex, neuronal types defined by scRNA-seq may not map uniquely onto the types derived from electrophysiology and/or morphology.
42. Cadwell CR, Palasantza A, Jiang X, Berens P, Deng Q, Yilmaz M, Reimer J, Shen S, Bethge M, Tolias KF *et al.*: **Electrophysiological, transcriptomic and morphologic profiling of single neurons using Patch-seq.** *Nat Biotechnol* 2016, **34**:199-203.
 43. Földy C, Darmanis S, Aoto J, Malenka RC, Quake SR, Südhof TC: **Single-cell RNAseq reveals cell adhesion molecule profiles in electrophysiologically defined neurons.** *Proc Natl Acad Sci* 2016, **113**:E5222-E5231.
 44. Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, Ailamaki A, Alonso-Nanclares L, Antille N, Arsever S *et al.*: **Reconstruction and simulation of neocortical microcircuitry.** *Cell* 2015, **163**:456-492.
 45. Klein AM, Mazutis L, Akartuna I, Tallapragada N, Veres A, Li V, Peshkin L, Weitz DA, Kirschner MW: **Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells.** *Cell* 2015, **161**:1187-1201.
 46. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR, Kamitaki N, Martersteck EM *et al.*: **Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets.** *Cell* 2015, **161**:1202-1214.
- Droplet-based scRNA-seq developed by this group and by Klein *et al.* [45] expands throughput by orders of magnitude, enabling higher sample sizes needed for heterogeneous tissues like cortex. Here, a novel analysis in retina also provides a template for unbiased sampling of tissues that contain both highly abundant and very rare cell types of interest.
47. Masland RH: **The neuronal organization of the retina.** *Neuron* 2012, **76**:266-280.
 48. Telley L, Govindan S, Prados J, Stevant I, Nef S, Dermitzakis E, Dayer A, Jabaudon D: **Sequential transcriptional waves direct the differentiation of newborn neurons in the mouse neocortex.** *Science* 2016, **351**:1443-1446.
 49. Pollen AA, Nowakowski TJ, Chen J, Retallack H, Sandoval-Espinosa C, Nicholas CR, Shuga J, Liu SJ, Oldham MC, Diaz A *et al.*: **Molecular identity of human outer radial glia during cortical development.** *Cell* 2015, **163**:55-67.
- This study and Thomsen *et al.* [50] used distinct experimental designs to arrive at similar results searching for genes enriched in known subtypes of human fetal cortex progenitors. These authors performed extensive validations, and find important developmental differences that highlight the need for larger studies across more ages.

50. Thomsen ER, Mich JK, Yao Z, Hodge RD, Doyle AM, Jang S, Shehata SI, Nelson AM, Shapovalova NV, Levi BP *et al.*: **Fixed single-cell transcriptomic characterization of human radial glial diversity**. *Nat Methods* 2016, **13**:87-93.
- In addition to identifying a novel transcriptional signature of outer radial glial progenitors in human fetal cortex, this work presents a new method for isolating specific cell types of interest from non-genetic models, which will be critical for future scRNA-seq work in primates and other species.
51. Camp JG, Badsha F, Florio M, Kanton S, Gerber T, Wilsch-Bräuninger M, Lewitus E, Sykes A, Hevers W, Lancaster M *et al.*: **Human cerebral organoids recapitulate gene expression programs of fetal neocortex development**. *Proc Natl Acad Sci* 2015, **112**:15672-15677.
52. Stancik EK, Navarro-Quiroga I, Sellke R, Haydar TF: **Heterogeneity in ventricular zone neural precursors contributes to neuronal fate diversity in the postnatal neocortex**. *J Neurosci* 2010, **30**:7028-7036.
53. Hansen DV, Lui JH, Parker PRL, Kriegstein AR: **Neurogenic radial glia in the outer subventricular zone of human neocortex**. *Nature* 2010, **464**:554-561.
54. Fietz SA, Kelava I, Vogt J, Wilsch-Bräuninger M, Stenzel D, Fish JL, Corbeil D, Riehn A, Distler W, Nitsch R *et al.*: **OSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling**. *Nat Neurosci* 2010, **13**:690-699.
55. Betizeau M, Cortay V, Patti D, Pfister S, Gautier E, Bellemin-Ménard A, Afanassieff M, Huisoud C, Douglas RJ, Kennedy H *et al.*: **Precursor diversity and complexity of lineage relationships in the outer subventricular zone of the primate**. *Neuron* 2013, **80**:442-457.
56. Handel AE, Chintawar S, Lalic T, Whiteley E, Vowles J, Giustacchini A, Argoud K, Sopp P, Nakanishi M, Bowden R *et al.*: **Assessing similarity to primary tissue and cortical layer identity in induced pluripotent stem cell-derived cortical neurons through single-cell transcriptomics**. *Hum Mol Genet* 2016, **25**:989-1000.
57. Treutlein B, Lee QY, Camp JG, Mall M, Koh W, Shariati SAM, Sim S, Neff NF, Skotheim JM, Wernig M *et al.*: **Dissecting direct reprogramming from fibroblast to neuron using single-cell RNA-seq**. *Nature* 2016, **534**:391-395.
58. Klemm S, Semrau S, Wiebrands K, Mooijman D, Faddah DA, Jaenisch R, van Oudenaarden A: **Transcriptional profiling of cells sorted by RNA abundance**. *Nat Methods* 2014, **11**:549-551.
59. Pellegrino M, Sciambi A, Yates JL, Mast JD, Silver C, Eastburn DJ: **RNA-Seq following PCR-based sorting reveals rare cell transcriptional signatures**. *BMC Genomics* 2016, **17**:361.
60. Lee JH, Daugharthy ER, Scheiman J, Kalhor R, Yang JL, Ferrante TC, Terry R, Jeanty SSF, Li C, Amamoto R *et al.*: **Highly multiplexed subcellular RNA sequencing in situ**. *Science* 2014, **343**:1360-1363.
61. Satija R, Farrell JA, Gennert D, Schier AF, Regev A: **Spatial reconstruction of single-cell gene expression data**. *Nat Biotechnol* 2015, **33**:495-502.
62. Achim K, Pettit J-B, Saraiva LR, Gavriouchkina D, Larsson T, Arendt D, Marioni JC: **High-throughput spatial mapping of single-cell RNA-seq data to tissue of origin**. *Nat Biotechnol* 2015, **33**:503-509.
63. Macaulay IC, Haerty W, Kumar P, Li YI, Hu TX, Teng MJ, Goolam M, Saurat N, Coupland P, Shirley LM *et al.*: **G&T-seq: parallel sequencing of single-cell genomes and transcriptomes**. *Nat Methods* 2015, **12**:519-522.
64. Dey SS, Kester L, Spanjaard B, Bienko M, van Oudenaarden A: **Integrated genome and transcriptome sequencing of the same cell**. *Nat Biotechnol* 2015, **33**:285-289.
65. Kharchenko PV, Silberstein L, Scadden DT: **Bayesian approach to single-cell differential expression analysis**. *Nat Methods* 2014, **11**:740-742.
66. Fan J, Salathia N, Liu R, Kaeser GE, Yung YC, Herman JL, Kaper F, Fan J-B, Zhang K, Chun J *et al.*: **Characterizing transcriptional heterogeneity through pathway and gene set overdispersion analysis**. *Nat Methods* 2016, **13**:241-244.